

H^1 - L^1 Boundedness of Rough Toroidal Pseudo-Differential Operator

Ramla Benhamoud

School of Mathematical Sciences, Zhejiang Normal University, Jinhua, China

ABSTRACT

In this paper, we study rough toroidal pseudo-differential operators. We prove that they are bounded from H^1 to L^1 on torus for symbols in a critical rough toroidal class.

KEYWORDS: Rough toroidal pseudo-differential operator; Rough toroidal symbol; H^1 (T^n); Torus; Atomic decomposition

Mathematics Subject Classification

Primary: 42B05,
Secondary: 35S05.

How to cite this paper: Ramla Benhamoud "H1-L1 Boundedness of Rough Toroidal Pseudo-Differential Operator" Published in International Journal of Trend in Scientific Research and Development (ijtsrd), ISSN: 2456-6470, Volume-8 | Issue-1, February 2024, pp.263-270, URL: www.ijtsrd.com/papers/ijtsrd62379.pdf



IJTSRD62379

Copyright © 2024 by author (s) and International Journal of Trend in Scientific Research and Development Journal. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0) (<http://creativecommons.org/licenses/by/4.0>)



INTRODUCTION

The theory of pseudo-differential operators was started by Kohn and Nirenberg and Hörmander. These operators have wide use in the study of partial differential equations and harmonic analysis. They are defined on \mathbb{R}^n as follows

$$T_a f(x) = \int_{\mathbb{R}^n} e^{ix\xi} a(x, \xi) \hat{f}(\xi) d\xi,$$

where \hat{f} is the Fourier transform of f and $a(x, \xi)$ is called the symbol of the operator T_a . According to the behavior of symbols and their derivatives, there are different symbol classes.

Let \mathbb{N} be the set $\{1, 2, 3, \dots\}$ and \mathbb{N}_0 the set $\{0, 1, 2, \dots\}$. We recall the classical symbol class introduced in which is known by Hörmander class $S_{\rho, \delta}^m(\mathbb{R}^n \times \mathbb{R}^n)$ ($m \in \mathbb{R}$ and $0 \leq \rho, \delta \leq 1$). It is the set of functions a that are smooth in (x, ξ) and satisfy

$$|\nabla_{\xi}^N \nabla_x^M a(x, \xi)| \leq C(1 + |\xi|)^{m - \rho N + \delta M}$$

for any $N, M \in \mathbb{N}_0$.

The concept of periodic pseudo-differential operators was introduced by Agranovich. He presented a global quantization on the unit circle S^1 that could be

extended to the torus T^n . These operators can be associated with the symbols $a: T^n \times \mathbb{R}^n \mapsto \mathbb{C}$ or $a: T^n \times \mathbb{R}^n \mapsto \mathbb{C}$ and denoted by $a(x, D)$. The equivalence between them has been proven in .

Let the torus $T^n = (\mathbb{R}/2\pi\mathbb{Z})^n$ and $\mathcal{S}(\mathbb{Z}^n)$ be the restriction of schwartz space $\mathcal{S}(\mathbb{R}^n)$ on \mathbb{Z}^n . We define the toroidal Fourier transform $\mathcal{F}_{T^n}: C^\infty(T^n) \rightarrow \mathcal{S}(\mathbb{Z}^n)$ by

$$(\mathcal{F}_{T^n} f)(\xi) = \hat{f}(\xi) = (2\pi)^{-n} \int_{T^n} e^{-ix\xi} f(x) dx.$$

Then, \mathcal{F}_{T^n} is a bijection and its inverse $\mathcal{F}_{T^n}^{-1}: \mathcal{S}(\mathbb{Z}^n) \rightarrow C^\infty(T^n)$ is given by

$$f(x) = \sum_{\xi \in \mathbb{Z}^n} e^{ix\xi} (\mathcal{F}_{T^n} f)(\xi) = \sum_{\xi \in \mathbb{Z}^n} e^{ix\xi} \hat{f}(\xi).$$

In , Cardona et al. defined the periodic Hörmander class $S_{\rho, \delta}^m(T^n \times \mathbb{R}^n)$ as a class of Hörmander symbols $S_{\rho, \delta}^m(\mathbb{R}^n \times \mathbb{R}^n)$ which are 2π -periodic in x , i.e., the set of smooth functions in $(x, \xi) \in T^n \times \mathbb{R}^n$ and satisfy

$$|\partial_x^\beta \partial_\xi^\alpha a(x, \xi)| \leq C_{\alpha, \beta} \langle \xi \rangle^{m - \rho|\alpha| + \delta|\beta|}.$$

The corresponding periodic pseudo-differential operator is given by

$$a(x, D)f(x) = (2\pi)^{-n} \int_{\mathbb{T}^n} \int_{\mathbb{R}^n} e^{i(x-y)\xi} a(x, \xi) f(y) d\xi dy.$$

Let Δ_{ξ_j} , $\xi \in \mathbb{Z}^n$, $j = 1, \dots, n$ be the forward partial difference operators such that for each function $f: \mathbb{Z}^n \rightarrow \mathbb{C}$

$$\Delta_{\xi_j} f(\xi) = f(\xi + \delta_j) - f(\xi),$$

and for $\alpha \in \mathbb{N}_0^n$

$$\Delta_{\xi}^{\alpha} = \Delta_{\xi_1}^{\alpha_1} \dots \Delta_{\xi_n}^{\alpha_n}.$$

The toroidal class $S_{\rho, \delta}^m(\mathbb{T}^n \times \mathbb{Z}^n)$, $0 \leq \rho, \delta \leq 1$ is the set of functions $a(x, \xi)$ that are smooth in x for all $\xi \in \mathbb{Z}^n$ and satisfy

$$|\Delta_{\xi}^{\alpha} \partial_x^{\beta} a(x, \xi)| \leq C_{\alpha, \beta} \langle \xi \rangle^{m - \rho|\alpha| + \delta|\beta|}$$

for all $\alpha, \beta \in \mathbb{N}_0^n$. The toroidal pseudo-differential operators are defined by

$$a(x, D)f(x) = \sum_{\xi \in \mathbb{Z}^n} e^{ix\xi} a(x, \xi) \hat{f}(\xi), \quad f \in C^{\infty}(\mathbb{T}^n).$$

Turunen and Vainikko studied products and asymptotic expansions for adjoints of periodic pseudo-differential operators. More results about the calculations of these operators have been presented by Ruzhansky and Turunen in [1] and [2].

The boundedness of pseudo-differential operators for symbols in Hörmander class $S_{\rho, \delta}^m$ is an interesting topic that has been extensively studied in the theory of these operators. We refer to [3] for results on \mathbb{R}^n . Many studies have been conducted to explore the boundedness of periodic pseudo-differential operators on torus \mathbb{T}^n and Lie groups. For instance, when symbols with finite regularity in $x \in \mathbb{T}^n$, Ruzhansky and Turunen proved the L^2 boundedness for $|\partial_x^{\beta} a(x, \xi)| \leq C$ for all $(x, \xi) \in \mathbb{T}^n \times \mathbb{Z}^n$ and $|\beta| \leq n + 1$ and without any regularity require for the variable ξ which is better than results on \mathbb{R}^n since the torus \mathbb{T}^n is compact. In [4], the L^2 boundedness has been shown for $a \in S_{\rho, \delta}^0(\mathbb{T}^n \times \mathbb{Z}^n)$ ($0 \leq \delta < \rho \leq 1$). The same result has been proved by Fischer [5] for the compact Lie group G . Delagdo [6] studied L^p boundedness, he showed that for $a \in S_{\rho, 0}^{n/2(\rho-1)}(\mathbb{T}^n \times \mathbb{Z}^n)$, $0 < \rho < 1$, the periodic pseudo-differential operator is bounded from L^{∞} into BMO , and by interpolation he concluded L^p ($2 \leq p < \infty$) boundedness. This result was extended to compact Lie group G in [7]. Delagdo and Ruzhansky extended also L^p boundedness for $1 < p < 2$ by using duality. They proved also for symbols with finite regularity that the operator $a(x, D)$ is bounded on $L^p(\mathbb{T}^n)$ ($1 < p < \infty$) for $S_{\rho, \delta}^m(\mathbb{T}^n \times \mathbb{Z}^n)$, $m = -\left[\frac{n}{2}\right](1 - \rho)|1/p - 1/2| - \delta\left(\left[\frac{n}{p}\right] + 1\right)$ and $|\beta| \leq \left[\frac{n}{p}\right] + 1$, where $0 \leq \delta < \rho < 1$. For more results concerning $L^p(\mathbb{T}^n)$ ($1 < p < \infty$) boundedness, one can see [8].

The weak type (1,1) was studied in [9], Cardona proved that for $a \in S_{1, \delta}^0(\mathbb{T}^n \times \mathbb{Z}^n)$, $0 \leq \delta < 1$, the pseudo-differential operator is weak type (1,1) on torus \mathbb{T}^n . For the multi-linear pseudo-differential operators on $\mathbb{T}^n \times \mathbb{Z}^n$, we refer to [10] and references therein. Furthermore, in [11], Cardona et al. showed that the boundedness of Fourier integral operators in $L^p(\mathbb{R}^n)$ ($1 < p < \infty$) implies the boundedness of periodic Fourier integral operators in $L^p(\mathbb{T}^n)$ ($1 < p < \infty$), i.e., for pseudo-differential operators is also true.

In [12], Kenig and Staubach defined a rough Hörmander symbol class $L^{\infty} S_{\rho}^m$ which symbols behave in the spatial variable like an L^{∞} function. It is known that $S_{\rho, 1}^m \subset L^{\infty} S_{\rho}^m$. Recently, Guo and Zhu studied endpoint estimates for the symbol class $a \in S_{\rho, 1}^{n(\rho-1)}(\mathbb{R}^n \times \mathbb{R}^n)$ and rough class $L^{\infty} S_{\rho}^{n(\rho-1)}$. We recall their Theorem 1.2.

Theorem A 1. *Let $a \in L^{\infty} S_{\rho}^{n(\rho-1)}(\mathbb{R}^n \times \mathbb{R}^n)$, $0 \leq \rho < 1$. Then the operator T_a is bounded from $H^1(\mathbb{R}^n)$ to $L^1(\mathbb{R}^n)$.*

For $\rho = 1$ they constructed a symbol $a \in S_{1, 1}^0$ such that T_a is unbounded from H^1 to L^1 . Similarly, we state the definition of rough toroidal symbol class $L^{\infty} S_{\rho}^m(\mathbb{T}^n \times \mathbb{Z}^n)$.

Definition 1. *The class $L^{\infty} S_{\rho}^m(\mathbb{T}^n \times \mathbb{Z}^n)$, $0 \leq \rho \leq 1$ is the set of functions which are measurable in x and satisfy $\|\Delta_{\xi}^{\alpha} a(\cdot, \xi)\|_{L^{\infty}(\mathbb{T}^n)} \lesssim \langle \xi \rangle^{m - \rho|\alpha|}$.*

In this note, we consider the rough toroidal class $L^{\infty} S_{\rho}^m(\mathbb{T}^n \times \mathbb{Z}^n)$ and we obtain a periodic version of Theorem A. Our main result is the following:

Theorem 1. *If $a \in L^{\infty} S_{\rho}^{n(\rho-1)}(\mathbb{T}^n \times \mathbb{Z}^n)$, $0 \leq \rho < 1$, then the toroidal pseudo-differential operator $a(x, D)$ is bounded from $H^1(\mathbb{T}^n)$ to $L^1(\mathbb{T}^n)$.*

Remark 1. *Theorem 1 extends the symbol class where the pseudo-differential operators are $H^1 - L^1$ bounded on torus to $S^{\wedge\{n(\rho-1)\}}_{\rho, \delta}$ and $0 \leq \delta \leq 1$. Our proof is based on the proprieties of the difference operator, i.e., we follow the method used in [13] by using the difference operator instead of the derivative.*

This note is organized as follows. In Section 2, we recall some basic notions and lemmas about the periodic analysis of pseudo-differential operators and the atomic characterization of Hardy spaces on a torus \mathbb{T}^n . In Section 3, we prove Theorem 1.

Preliminary notions and lemmas

The following lemma is a periodic version of Proposition 2.3 in [14].

Lemma 1. Assume $0 \leq \rho \leq 1$, $a \in L^\infty S_\rho^{n(\rho-1)}(\mathbb{T}^n \times \mathbb{Z}^n)$ and $1 < p < \infty$. Then the operator $a(x, D)$ is bounded from $L^p(\mathbb{T}^n)$ to $L^p(\mathbb{T}^n)$.

Proof. Since from , the $L^p(1 < p < \infty)$ boundedness of pseudo-differential operators on \mathbb{R}^n implies the boundedness of periodic operators on \mathbb{T}^n . Then we get directly Lemma 1 from. \square

For two multi-indices $\alpha, \beta \in \mathbb{N}_0^n$, $\beta \leq \alpha$ means that $\beta_k \leq \alpha_k$ for all $1 \leq k \leq n$ and we denote

$$\binom{\alpha}{\beta} = \prod_{k=1}^n \binom{\alpha_k}{\beta_k} = \prod_{k=1}^n \frac{\alpha_k!}{(\alpha_k - \beta_k)! \beta_k!}.$$

We recall some known proprieties for the difference operator Δ_ξ .

Lemma 2. (i) Let $f, g: \mathbb{Z}^n \rightarrow \mathbb{C}$. There holds $\Delta_\xi^\alpha(fg)(\xi) = \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \Delta_\xi^\beta f \Delta_\xi^{\alpha-\beta} g(\xi + \beta)$.

Lemma 3. Let $f, g: \mathbb{Z}^n \rightarrow \mathbb{C}$. Then $\sum_{\xi \in \mathbb{Z}^n} f(\xi) [\Delta_\xi^\gamma g](\xi) = (-1)^\gamma \sum_{\xi \in \mathbb{Z}^n} [\bar{\Delta}_\xi^\gamma f](\xi) g(\xi)$, where $\bar{\Delta}_\xi^\gamma f(\xi) = f(\xi) - f(\xi - \delta_j)$.

Atomic decomposition for $H^p(\mathbb{T}^n)$:

As to the Euclidean case \mathbb{R}^n , Hardy space on a torus $\mathbb{T}^n (n \geq 1)$ admits atomic decomposition. For more details, we refer to and , where Folland and Stein showed atomic decomposition for Hardy spaces on homogeneous groups and in Qing et al. for Hardy spaces on graded Lie groups. And for $n = 1$, we can see .

$$\begin{aligned} \sum_{\xi \in \mathbb{Z}^n - \{0\}} |\hat{b}(\xi)| |\xi|^{-n} &= \sum_{|\xi| \leq [r^{-1}]} |\hat{b}(\xi)| |\xi|^{-n} + \sum_{|\xi| > [r^{-1}]} |\hat{b}(\xi)| |\xi|^{-n} \\ &\lesssim \sum_{|\xi| \leq [r^{-1}]} |\xi|^{-n} \int_{B(x_0, r)} |e^{-ix\xi} - 1| |b(x)| dx + \sum_{|\xi| > [r^{-1}]} |\hat{b}(\xi)| |\xi|^{-n} \\ &\lesssim \sum_{|\xi| \leq [r^{-1}]} |\xi|^{\theta-n} \int_{B(x_0, r)} |x|^\theta |b(x)| dx + \left(\sum_{|\xi| > [r^{-1}]} |\hat{b}(\xi)|^2 \right)^{1/2} \left(\sum_{|\xi| > [r^{-1}]} |\xi|^{-2n} \right)^{1/2} \\ &\lesssim \sum_{|\xi| \leq [r^{-1}]} |\xi|^{\theta-n} \|b\|_{L^2} \left(\int_{B(x_0, r)} |x|^{2\theta} dx \right)^{1/2} + \|b\|_{L^2} \left(\sum_{|\xi| > [r^{-1}]} |\xi|^{-2n} \right)^{1/2} \\ &\lesssim \sum_{|\xi| \leq [r^{-1}]} |\xi|^{\theta-n} r^{-n/2} r^{3n\theta/2} + r^{-n/2} \left(\sum_{|\xi| > [r^{-1}]} |\xi|^{-2n} \right)^{1/2} \\ &\lesssim \sum_{|\xi| \leq [r^{-1}]} |\xi|^{\theta-n} r^{-n/2} r^{3n\theta/2} + r^{-n/2} r^{n/2} \left(\sum_{|\xi| > [r^{-1}]} |\xi|^{-n} \right)^{1/2} \\ &\lesssim 1 \end{aligned}$$

we can take $\theta = 1/3$.

Case 2. If $r \geq 1$ we can get

$$\sum_{\xi \in \mathbb{Z}^n - \{0\}} |\hat{b}(\xi)| |\xi|^{-n} \leq \left(\sum_{\xi \in \mathbb{Z}^n - \{0\}} |\hat{b}(\xi)|^2 \right)^{1/2} \left(\sum_{\xi \in \mathbb{Z}^n - \{0\}} |\xi|^{-2n} \right)^{1/2} \lesssim 1.$$

Definition 2. Let $0 < p \leq 1 \leq q \leq \infty$, $p \wedge q \neq 1$ and s is an integer at least $[n(1/p - 1)]$. We say a function b on \mathbb{T}^n is a (p, q, s) -atom if it is a compactly supported L^q function such that

- there is a ball B such that $\text{supp } b \subset \bar{B}$ and $\|b\|_{L^q(B)} \leq |B|^{1/q-1/p}$.
- $\int_{\mathbb{T}^n} b(x) x^\alpha dx = 0$, where $0 \leq |\alpha| = \alpha_1 + \alpha_2 + \dots + \alpha_n \leq s$ and $x^\alpha = x_1^{\alpha_1} x_2^{\alpha_2} \dots x_n^{\alpha_n}$.

The atomic decomposition of $H^p(\mathbb{T}^n)$ is given as follows.

Proposition 1. Let (p, q, s) be as in Definition 2. Then for all any (p, q, s) atom a , one has $\|a\|_{H^p(\mathbb{T}^n)} \lesssim 1$. Conversely, for any $f \in H^p(\mathbb{T}^n)$, there exists a sequence $\{\lambda_j\}_{j=1}^\infty$ such that $f = \sum_{j=1}^\infty \lambda_j a_j$ with convergence in $\mathcal{S}'(\mathbb{T}^n)$ and $(\sum_{j=1}^\infty |\lambda_j|^p)^{1/p} \lesssim \|f\|_{H^p(\mathbb{T}^n)}$.

Before giving the proof of our main result, we present the following inequality which will be used in our proof.

Lemma 4. For any L^2 -atom b for $H^1(\mathbb{T}^n)$ and $n \geq 1$, we have, $\sum_{\xi \in \mathbb{Z}^n - \{0\}} |\hat{b}(\xi)| |\xi|^{-n} \leq C$.

Proof. From Definition 2, we have $\text{supp } b \subset B(x_0, r)$, $\int_{B(x_0, r)} b(x) dx = 0$ and $\|b\|_{L^2} \leq r^{-n/2}$.

Case 1. If $r < 1$, by using Hölder inequality and the inequality in $|e^{-ix\xi} - 1| \lesssim |x|^\theta |\xi|^\theta$ which holds for all $\theta \in (0, 1)$, we can get

Proof of our main result

To start proving Theorem 1, we use atomic decomposition for $H^1(\mathbb{T}^n)$. It is enough to show $\|a(x, D)b\|_{L^1(\mathbb{T}^n)} \lesssim 1$ for a L^2 -atom b . We prove the case $\rho = 0$ latter, since the kernel is not rapidly decreasing. First, we consider $0 < \rho < 1$.

Case of $0 < \rho < 1$

We denote the kernel of $a(x, D)$ by

$$k(x, y) = \sum_{\xi \in \mathbb{Z}^n} e^{i(x-y)\xi} a(x, \xi).$$

From summation by parts in Lemma 3, one can easily show the equality

$$(e^{-i(x-y)} - 1)^\gamma e^{i(x-y)\xi} = (-1)^{|\gamma|} \bar{\Delta}_\xi^\gamma e^{i(x-y)\xi},$$

and since we are working on the torus \mathbb{T}^n , it is easy to check that the following inequality in

$$|x - y|^N \leq c \sum_{|\gamma|=N} |(e^{-i(x-y)} - 1)^\gamma|,$$

is true for all $x, y \in \mathbb{T}^n$. From [3.1] and [3.2], it yields

$$\begin{aligned} (e^{-i(x-y)} - 1)^\gamma k(x, y) &= \sum_{\xi \in \mathbb{Z}^n} (e^{-i(x-y)} - 1)^\gamma e^{i\xi(x-y)} a(x, \xi) \\ &= \sum_{\xi \in \mathbb{Z}^n} (-1)^{|\gamma|} [\bar{\Delta}_\xi^\gamma e^{i\xi(x-y)}] a(x, \xi) \\ &= \sum_{\xi \in \mathbb{Z}^n} e^{i\xi(x-y)} \bar{\Delta}_\xi^\gamma [a(x, \xi)]. \end{aligned}$$

On the other hand, let $B_j = \{\xi \in \mathbb{Z}^n : 2^{j-1} \leq |\xi| \leq 2^{j+1}\}$ and $|\xi|_\infty = \max_i |\xi_i|$ where ξ_i represents the i -th component of the vector $\xi \in \mathbb{Z}^n$. It is clear that $\mathbb{Z}^n = \bigcup_{j \in \mathbb{N}_0} B_j$. If we choose $\mu = 2^{j+1}$ and $0 < \epsilon < 1$, we can write

$$\begin{aligned} \sum_{\xi \in B_j} |\xi|^{\epsilon-n} &\lesssim \sum_{l \in \mu^{-1}\mathbb{Z}^n \setminus \{0\}, |l|_\infty \leq 1} |\mu l|^{\epsilon-n} \\ &\lesssim \sum_{\substack{\xi \in \mathbb{Z}^n \setminus \{0\}, |\xi|_\infty \leq \mu \\ [\mu]+1}} |\xi|^{\epsilon-n} \\ &\lesssim \sum_{N=1}^{\mu} \sum_{\xi \in Y_N} |\xi|^{\epsilon-n}, \end{aligned}$$

such that

$$\begin{aligned} \mathbb{Z}^n \setminus \{0\} &= \bigcup_{N \in \mathbb{N}} \{\xi \in \mathbb{Z}^n : |\xi|_\infty = N\} \\ &= \bigcup_{N \in \mathbb{N}} Y_N \end{aligned}$$

and $\text{card } Y_N \leq CN^{n-1}$. Then, we can obtain

$$\begin{aligned} \sum_{\xi \in B_j} |\xi|^{\epsilon-n} &\lesssim \sum_{N=1}^{[\mu]+1} N^{n-1} N^{\epsilon-n} \\ &\lesssim \int_0^{[\mu]} x^{\epsilon-1} dx \\ &\lesssim 2^{j\epsilon}. \end{aligned}$$

So, by some simple computations, we can estimate the kernel $k(x, y)$ as follows

$$\begin{aligned}
 |k(x, y)| &\lesssim |e^{i(x-y)} - 1|^{-N} \sum_{\xi \in \mathbb{Z}^n} |\Delta_\xi^N a(x, \xi)| \\
 &\lesssim |x - y|^{-N} \sum_{\xi \in \mathbb{Z}^n} \langle \xi \rangle^{n(\rho-1)-N\rho} \\
 &\lesssim |x - y|^{-N} \sum_{j \in \mathbb{N}_0} \sum_{\xi \in B_j} \langle \xi \rangle^{n(\rho-1)-N\rho} \\
 &\lesssim |x - y|^{-N} \sum_{j \in \mathbb{N}_0} 2^{jn(\rho-1)-jN\rho+jn-j\epsilon} \sum_{\xi \in B_j} \langle \xi \rangle^{\epsilon-n} \\
 &\lesssim |x - y|^{-N} \sum_{j \in \mathbb{N}_0} 2^{jn\rho-jN\rho} \\
 &\lesssim |x - y|^{-N}
 \end{aligned}$$

the last inequality is true for $N > n$. Then, if $|x| > 2r$, it yields

$$|a(x, D)b(x)| = \left| \int_{|y| \leq r} k(x, y)b(y)dy \right| \lesssim \int_{|y| \leq r} |x - y|^{-N}|b(y)|dy \lesssim |x|^{-N}.$$

Part 1

Assume $r \geq 1$, then

$$\begin{aligned}
 \|a(x, D)b\|_{L^1(\mathbb{T}^n)} &= \int_{\mathbb{T}^n \setminus B(0, 2r)} |a(x, D)b(x)|dx + \int_{B(0, 2r)} |a(x, D)b(x)|dx \\
 &\lesssim \int_{\mathbb{T}^n \setminus B(0, 2r)} |x|^{-N}dx + r^{n/2} \|a(x, D)b\|_2 \\
 &\lesssim 1 + r^{n/2} \|b\|_2 \lesssim 1.
 \end{aligned}$$

Part 2

For $r < 1$, we use discrete dyadic decomposition $(\varphi_j)_{j \in \mathbb{N}_0} \subset \mathcal{S}(\mathbb{Z}^n)$ as in .

- $\text{supp } \varphi_0 \subset \{\xi \in \mathbb{Z}^n: |\xi| \leq 2\}$ and $\varphi_0(\xi) = 1$ when $|\xi| \leq 1$.
- $\text{supp } \varphi_j \subset \{\xi \in \mathbb{Z}^n: 2^{j-1} < |\xi| \leq 2^{j+1}\}$ for $j \geq 1$.
- For any $\xi \in \mathbb{Z}^n$, one have $0 \leq \varphi_j(\xi) \leq 1$ ($j \in \mathbb{N}_0$) and $\sum_{j \in \mathbb{N}_0} \varphi_j(\xi) = 1$.
- For any $\alpha \in \mathbb{N}_0^n$, there exists a constant $C_\alpha > 0$ such that

$$|\Delta_\xi^\alpha \varphi_j(\xi)| \leq C_\alpha \langle \xi \rangle^{-|\alpha|}$$

for any $j \in \mathbb{N}$, $\xi \in \mathbb{Z}^n$.

Then, we can write

$$\begin{aligned}
 a(x, D)b(x) &= \sum_{j \in \mathbb{N}_0} \sum_{\xi \in B_j} \int_{\mathbb{T}^n} e^{i(x-y)\xi} a(x, \xi) \varphi_j(\xi) b(y) dy \\
 &= \sum_{j \in \mathbb{N}_0} a_j(x, D)b(x),
 \end{aligned}$$

where $B_j = \{\xi \in \mathbb{Z}^n: 2^{j-1} \leq |\xi| \leq 2^{j+1}\}$ and $B_0 = \{\xi \in \mathbb{Z}^n: |\xi| \leq 2\}$.

Using the L^2 boundedness result in Lemma 1, we can get

$$\int_{|x| \leq 3r} \left| \sum_{j \in \mathbb{N}_0} a_j(x, D)b(x) \right| dx \lesssim 1.$$

Now we shall prove

$$\int_{|x| > 3r} |a_j(x, D)b(x)| dx \lesssim 1.$$

Assume $2^j \geq r^{-1/\rho}$, one can get

$$\begin{aligned}
 \sum_{2^j \geq r^{-1/\rho}} \int_{|x| > 3r} |a_j(x, D)b(x)| dx &\lesssim \sum_{2^j \geq r^{-1/\rho}} \int_{|x| > 3r} \int_{|y| \leq r} |e^{i(x-y)} - 1|^{-N} 2^{j\rho(n-N)} |b(y)| dy dx \\
 &\lesssim \sum_{2^j \geq r^{-1/\rho}} \int_{|x| > 3r} 2^{j\rho(n-N)} \int_{|y| \leq r} |x-y|^{-N} |b(y)| dy dx \\
 &\lesssim \sum_{2^j \geq r^{-1/\rho}} \int_{|x| > 3r} 2^{j\rho(n-N)} |x|^{-N} dx \\
 &\lesssim 1.
 \end{aligned}$$

For $2^j < r^{-1/\rho}$, we have

$$\sum_{2^j < r^{-1/\rho}} \int_{|x| > 3r} |a_j(x, D)b(x)| dx \lesssim \sum_{2^j < r^{-1/\rho}} \int_{|x| > 3r} \sum_{\xi \in B_j} |x|^{-N} |\Delta_\xi^N [a(x, \xi) \varphi_j(\xi) \hat{b}(\xi)]|.$$

From discrete Leibniz formula Lemma 2, and for $\alpha \neq N$, we can obtain

$$\begin{aligned}
 |\Delta_\xi^{N-\alpha} \hat{b}(\xi + \alpha)| &= |\Delta_\xi^{N-\alpha} \left(\int_{\mathbb{T}^n} e^{-iy(\xi+\alpha)} b(y) dy \right)| \\
 &\leq \int_{\mathbb{T}^n} |y|^{N-\alpha} |b(y)| dy \\
 &\leq \int_{|y| \leq r} r^{N-\alpha} |b(y)| dy \\
 &\lesssim r^{N-\alpha}.
 \end{aligned}$$

Then, we can get the following estimate

$$\begin{aligned}
 &|\Delta_\xi^N [a(x, \xi) \varphi_j(\xi) \hat{b}(\xi)]| \\
 &\leq \sum_{0 \leq \alpha \leq N} |\Delta_\xi^\alpha [a(x, \xi) \varphi_j(\xi)]| |\Delta_\xi^{N-\alpha} \hat{b}(\xi + \alpha)| \\
 &= |\Delta_\xi^N [a(x, \xi) \varphi_j(\xi)]| |\hat{b}(\xi + N)| + \sum_{0 \leq \alpha < N} 2^{jn(\rho-1)-\rho\alpha} r^{N-\alpha} \\
 &\lesssim 2^{jn(\rho-1)-j\rho N} |\hat{b}(\xi + N)| + \sum_{0 \leq \alpha < N} 2^{jn(\rho-1)} r^N (2^{-j\rho} r^{-1})^\alpha \\
 &\lesssim 2^{jn(\rho-1)-j\rho N} |\hat{b}(\xi + N)| + 2^{jn(\rho-1)} 2^{-j\rho N} 2^{j\rho} r \\
 &\lesssim 2^{jn(\rho-1)-j\rho N} [|\hat{b}(\xi + N)| + 2^{j\rho} r].
 \end{aligned}$$

So, from Lemma 4 we obtain

$$\begin{aligned}
 &\sum_{2^j < r^{-1/\rho}} \int_{|x| > 3r} |a_j(x, D)b(x)| dx \\
 &\leq \sum_{2^j < r^{-1/\rho}} \int_{|x| > 3r} \sum_{\xi \in B_j} |x|^{-N} 2^{jn(\rho-1)-j\rho N} [|\hat{b}(\xi + N)| + 2^{j\rho} r] dx \\
 &\lesssim \sum_{2^j < r^{-1/\rho}} \sum_{\xi \in B_j} 2^{jn(\rho-1)-j\rho N} [|\hat{b}(\xi + N)| + 2^{j\rho} r] \\
 &\lesssim \sum_{2^j < r^{-1/\rho}} \sum_{\xi \in B_j} 2^{jn(\rho-1)-j\rho N} |\hat{b}(\xi + N)| + \sum_{2^j < r^{-1/\rho}} \sum_{\xi \in B_j} 2^{jn(\rho-1)-j\rho N} 2^{j\rho} r \\
 &\lesssim \sum_{2^j < r^{-1/\rho}} \sum_{\xi \in B_j} 2^{jn(\rho-1)-j\rho N} |\hat{b}(\xi + N)| + \sum_{2^j < r^{-1/\rho}} 2^{j\rho(n-N)} 2^{j\rho} r \\
 &\lesssim \sum_{2^j < r^{-1/\rho}} \sum_{\xi \in B_j} 2^{j\rho n-j\rho N} |\hat{b}(\xi + N)| |\xi|^{-n} + \sum_{2^j < r^{-1/\rho}} 2^{j\rho(n-N)} 2^{j\rho} r \\
 &\lesssim 1.
 \end{aligned}$$

Case of $\rho = 0$

In this part, we shall show that the operator $a(x, D)$ is bounded from H^1 to L^1 for the symbol $a \in L^\infty S_0^{-n}(\mathbb{T}^n \times \mathbb{Z}^n)$. As in the case $0 < \rho < 1$ we first get the estimate for $r \geq 1$ then $r < 1$.

Part 1. Suppose $r \geq 1$, if $x < 2r$, from Lemma 1 we can get

$$\|a(x, D)b\|_{L^1(B(x_0, 2r))} \lesssim r^{n/2} \|a(x, D)b\|_{L^2(\mathbb{T}^n)} \lesssim r^{n/2} \|b\|_{L^2(\mathbb{T}^n)} \lesssim 1.$$

For $\{x \in \mathbb{T}^n: |x| \geq 2r\}$, we use summation by parts in Lemma 2

$$\begin{aligned} |a(x, D)b(x)| &= |x|^{-n} \left| \sum_{\xi \in \mathbb{Z}^n} e^{ix\xi} \Delta_\xi^n (a(x, \xi) \widehat{b}(\xi)) \right| \\ &= |x|^{-n} \sum_{l=0}^n \left| \sum_{\xi \in \mathbb{Z}^n} e^{ix\xi} \Delta_\xi^{n-l} a(x, \xi) \widehat{y^l b}(\xi) \right| \\ &= |x|^{-n} \sum_{l=0}^n |a_l(x, D)[y^l b](x)| \end{aligned}$$

where $a_l(x, D)$ is a periodic pseudo-differential operator with symbol $\Delta_\xi^{n-l} a(x, \xi)$. It is obvious that $\Delta_\xi^{n-l} a(x, \xi) \in L^\infty S_0^{-n}(\mathbb{T}^n \times \mathbb{Z}^n)$. Since from Lemma 1, the operator $a_l(x, D)$ is bounded on $L^2(\mathbb{T}^n)$, one can obtain

$$\begin{aligned} \int_{\mathbb{T}^n \setminus B(x_0, 2r)} |a(x, D)b(x)| dx &\lesssim \sum_{l=0}^n \int_{\mathbb{T}^n \setminus B(x_0, 2r)} |x|^{-n} |a_l(x, D)(y^l b)(x)| dx \\ &\lesssim \sum_{l=0}^n r^{-n/2} \|y^l b\|_{L^2(\mathbb{T}^n)} \\ &\lesssim \sum_{l=0}^n r^{-n+l} \lesssim 1. \end{aligned}$$

Part 2. For $r < 1$, applying Lemma 4, it yields

$$|a(x, D)b(x)| = \sum_{\xi \in \mathbb{Z}^n} e^{ix\xi} a(x, \xi) \widehat{b}(\xi) d\xi \leq \sum_{\xi \in \mathbb{Z}^n} |\widehat{b}(\xi)| |\xi|^{-n} d\xi \lesssim 1.$$

Then, for x in $\{x \in \mathbb{T}^n: |x| \leq 2\}$ we can get

$$\int_{|x| \leq 2} |a(x, D)b(x)| dx \lesssim 1.$$

Now, we estimate the part where x in $S = \{x \in \mathbb{T}^n: |x| > 2\}$. As above when $r \geq 1$, we can write

$$\begin{aligned} |a(x, D)b(x)| &\lesssim |x|^{-2n} \sum_{l=0}^{2n} \left| \sum_{\xi \in \mathbb{Z}^n} e^{ix\xi} \Delta_\xi^{2n-l} a(x, \xi) \widehat{y^l b}(\xi) \right| \\ &= |x|^{-2n} \sum_{l=0}^{2n} |\tilde{a}_l(x, D)(y^l b)(x)|, \end{aligned}$$

where $\tilde{a}_l(x, D)$ is a pseudo-differential operator with the symbol $\Delta_\xi^{2n-l} a(x, \xi)$ belongs to $L^\infty S_0^{-n}$. From Lemma 1 we have that $\tilde{a}_l(x, D)$ is bounded from $L^{n/(n-1)}(\mathbb{T}^n)$ to $L^{n/(n-1)}(\mathbb{T}^n)$ if $n \geq 2$ and from $L^2(\mathbb{T}^n)$ to $L^2(\mathbb{T}^n)$ if $n = 1$. Then, we have

$$\begin{aligned} \int_S |a(x, D)b(x)| dx &\lesssim \sum_{l=0}^{2n} \int_S |x|^{-2n} |\tilde{a}_l(y^l b)(x)| dx \\ &\lesssim 1 + \sum_{l=1}^{2n} \|y^l b\|_{L^{n/(n-1)}(\mathbb{T}^n)} \\ &\lesssim 1 + \sum_{l=1}^{2n} r^{l+\frac{n}{2}-1} \|b\|_{L^2(\mathbb{T}^n)} \lesssim 1. \end{aligned}$$

We use same computation for $n = 1$ by using L^2 boundedness.

The proof is completed.

Acknowledgments

Benhamoud Ramla was supported by the National Natural Science Foundation of China (No. 12071437).

Declarations

Conflict of interest The authors declare no competing interests.

References

- [1] Agranovich M. S.: Spectral properties of elliptic pseudodifferential operators on a closed curve. *Funktsional. Anal. i Prilozhen.* 13(4), 54-56 (1979)
- [2] Álvarez J., Hounie J.: Estimates for the kernel and continuity properties of pseudo-differential operators. *Ark Mat.* 28(1), 1-22 (1990)
- [3] Cardona D.: Weak type (1,1) bounds for a class of periodic pseudo-differential operators. *J. Pseudo-Differ. Oper. Appl.* 5(4), 507-515 (2014)
- [4] Cardona D.: On the boundedness of periodic pseudo-differential operators. *Monatsh. Math.* 185(2), 189-206 (2018)
- [5] Cardona D., Kumar V.: L^p -Boundedness and L^p -nuclearity of multilinear pseudo-differential operators on \mathbb{Z}^n and the torus \mathbb{T}^n . *J. Fourier Anal. Appl.* 25(6), 2973-3017 (2019)
- [6] Cardona D., Messiouene R., Senoussaoui A.: Periodic Fourier integral operators in L^p -spaces. *C. R. Math. Acad. Sci. Paris.* 359, 547-553 (2021)
- [7] Ching C. H.: Pseudo-differential operators with nonregular symbols. *J. Differential Equations* 11, 436-447 (1972)
- [8] Delgado J.: L^p -bounds for Pseudo-differential Operators on the Torus. *Oper. Theory Adv. Appl.* 231, 103-116 (2013)
- [9] Delgado J., Ruzhansky M.: L^p -bounds for pseudo-differential operators on compact Lie groups. *J. Inst. Math. Jussieu.* 18(3), 531-559 (2019)
- [10] Fischer V.: Intrinsic pseudo-differential calculi on any compact Lie group. *J. Funct. Anal.* 268(11), 3404-3477 (2015)
- [11] Folland G. B., Stein E. M.: Hardy spaces on homogeneous groups. Princeton University Press. 28, (1982)
- [12] Guo J., Zhu X.: Some notes on endpoint estimates for pseudo-differential operators. *Mediterr. J. Math.* 19(6), Paper No. 260, 14 pp (2022)
- [13] Hong Q., Hu G., Ruzhansky M.: Fourier multipliers for Hardy spaces on graded. Preprint 1-20 (2022) <https://arxiv.org/abs/2101.07499v2>
- [14] Hörmander L.: Pseudo-differential operators. *Comm. Pure Appl. Math.* 18, 501-517 (1965)
- [15] Hörmander L.: Pseudo-differential operators and hypoelliptic equations. *Singular integrals (Proc. Sympos. Pure Math., Vol. X, Chicago, Ill., 1966).* 138-183. Amer. Math. Soc., Providence, R.I., 1967
- [16] Hörmander L.: On the L^2 continuity of pseudo-differential operators. *Comm. Pure Appl. Math.* 24, 529-535 (1971)
- [17] Hounie J.: On the L^2 -continuity of pseudo-differential operators. *Comm. Partial Differential Equations.* 11(7), 765-778 (1986)
- [18] Kenig C. E., Staubach W.: ψ -pseudodifferential operators and estimates for maximal oscillatory integrals. *Studia Math.* 183(3), 249-258 (2007)
- [19] Kohn J. J., Nirenberg L.: An algebra of pseudo-differential operators. *Comm. Pure Appl. Math.* 18, 269-305 (1965)
- [20] Martínez B. B., Denk R., Monzón J. H., Nendel M.: Mapping properties for operator-valued pseudodifferential operators on toroidal besov spaces. *J. Pseudo-Differ. Oper. Appl.* 9(3), 523-538 (2018)
- [21] Mclean W.: Local and global description of periodic pseudo-differential operators. *Math. Nachr.* 150, 151-161 (1991)
- [22] Oswald P.: On Some Approximation Real Hardy Spaces Properties of real Hardy space ($0 < p \leq 1$). *J. Approx. Theory.* 40(1), 45-65 (1984)
- [23] Rodino L.: On the boundedness of pseudo differential operators in the class $L^m_{p,1}$. *Proc. Amer. Math. Soc.* 58, 211-215 (1976)
- [24] Ruzhansky M., Turunen V.: Quantization of Pseudo-differential Operators on the Torus. *J. Fourier Anal. Appl.* 16(6), 943-982 (2010)
- [25] Ruzhansky M., Turunen V.: Pseudo-differential operators and symmetries: background analysis and advanced topics. Birkhäuser-Verlag, Basel. (2010)
- [26] Stein E. M.: Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals, With the assistance of Timothy S. Murphy. Princeton Mathematical Series, 43. Monographs in Harmonic Analysis, III. Princeton University Press, Princeton, NJ, (1993)
- [27] Turunen V., Vainikko G.: On Symbol Analysis of Periodic Pseudo differential Operators. *Z. Anal. Anwendungen.* 17(1), 9-22 (1998)